Chemistry 210

Exam 2

Be sure to put your name on each page. This page can be removed from your exam so that you will have a Periodic Table handy throughout the exam, it does not need to be turned in. Show all your work for problems which require any sort of calculation, no credit will be given for answers without work shown. If you have shown a significant amount of work or multiple drawings for a problem, draw a box around what you consider your final answer.

Avogadro's Number = 6.022×10^{23} units/mol 32.00° F = 0.000° C = 273.15K Density of Water = $1.000^{g}/_{mL}$ $R = 0.08206^{\text{L*atm}}/_{\text{mol*K}} = 8.314^{\text{J}}/_{\text{mol*K}}$ PV=nRT $\Delta T_{fp/bp} = k_{fp/bp} \bullet m \bullet i$ For water, $k_{fp} = -1.86^{\circ C}/_{m}$; $k_{bp} = 0.52^{\circ C}/_{m}$ $\mathbf{P}_1 = \mathbf{X}_1 \mathbf{P}_1^{\,\mathbf{o}}$ P = cRTi $C_1V_1 = C_2V_2$ Quadratic formula: $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

$$k = Ae^{-Ea/RT}$$

$$ln(k) = \left(\frac{-E_a}{R}\right)\left(\frac{1}{T}\right) + ln(A)$$

$$ln\left(\frac{k_1}{k_2}\right) = \frac{E_a}{R}\left(\frac{1}{T_2} - \frac{1}{T_1}\right)$$

Integrated Rate Laws: 0th order $[A]_t = -kt + [A]_o$ 1st order $ln[A]_t = -kt + ln[A]_o$

58

Ce

140 12

90

Th

Pr

140 91

91

Pa

60

Nd

144 24

92

U

61

Pm

(145)

93

Np

62

Sm

150.36

94

Pu

63

Eu

151 97

95

Am (243)

64

Gd

157.25

96

Cm

65

Tb

158 93

97

Bk

Dy

162 50

98

Cf

2nd order

 $1/[A]_t = kt + 1/[A]_0$

\mathbf{H}																	He He
1.0079	4											5	6	7	8	9	4.0026
Li	Be											В	$\overset{\circ}{\mathbf{C}}$	N	Ŏ	F	Ne
6.941	9.0122											10.811	12.011	14.007	15.999	18.998	20.180
11	12											13	14	15	16	17	18
Na 22.990	Mg 24.305											Al 26.982	Si 28.086	P 30.974	S 32.066	Cl 35.453	Ar 39.948
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
39.098	40.078	44.956 39	47.88	50.942	51.996	54.938	55.847 44	58.933	58.69 46	63.546	65.39 48	69.723	72.61 50	74.922 51	78.96 52	79.904 53	83.80 54
Rb 85.468	Sr 87.62	Y 88.906	Zr	Nb 92,906	Mo 95.94	Tc	Ru	Rh	Pd 106.42	Ag	Cd	In 114.82	Sn	Sb	Te	I 126.90	Xe 131.29
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	T1	Pb	Bi	Po	At	Rn
132.91	137.33	138.91	178.49	180.95 105	183.84	186.21	190.23	192.22	195.08 110	196.97 111	200.59	204.38	207.2	208.98	(209)	(210)	(222)
Fr (223)	Ra	Ac 227.03	Rf (261)	Db (262)	Sg (263)	Bh (262)	Hs (265)	Mt (266)	(269)	(272)	(277)		117		110		

67

Ho

164.93

99

Es

68

Er

167.26

100

Fm

69

Tm

168 94

101

Md

71

Lu

174 97

103

Lr

(260)

Score

Yb

173 04

102

No

Multiple Choice: Circle the letter of the most correct response. (6pts. per question)

- 1. If the rate of a reaction increases by a factor if 4 when the initial concentration of reactant "A" is increased by a factor of 4, the reaction must be:
 - a. 1st order with respect to [A]o
 - b. 2nd order with respect to [A]o
 - c. The order of the reaction depends on the balanced chemical equation
 - d. 2nd order overall
 - e. 4th order with respect to [A]o
- 2. For a first order reaction:
 - a. The slope of the integrated rate law plot is equal to k
 - b. The intercept of the integrated rate law is equal to the *ln* of the initial concentration
 - c. The slope of the integrated rate law plot is equal to $(-E_a/R)$
 - d. The intercept of the integrated rate law plot is equal to the initial concentration
 - e. The slope of the integrated rate law is equal to the frequency factor, A.
- 3. Which of the following does *not* affect the rate of a reaction?
 - a. The frequency of collisions between reacting particles
 - b. The orientation of colliding particles
 - c. The energy of collisions between reacting particles
 - d. The coefficients of the reactants in the balanced equation
 - e. The temperature of the system
- 4. For the generic equation:

$$aA + bB \rightarrow cC + dD$$

Which of the following is a correct expression of the rate of the reaction:

- $\begin{array}{ll} a. & ^{-1}/_a \stackrel{\Delta[A]}{/_{\Delta t}} \\ b. & ^{-1}/_b \stackrel{\Delta[B]}{/_{\Delta t}} \end{array}$
- c. $k[A]^a[B]^b$ d. $\frac{1}{d}^{\Delta[D]}/_{\Delta t}$
- e. $k[C]^{c}[D]^{d}$
- 5. Which of the following is *true* regarding catalysts and catalyzed reactions?
 - a. The presence of a catalyst does not change the mechanism of a reaction
 - b. The concentration of a catalyst cannot appear in the rate law for a reaction
 - c. The presence of a catalyst changes the equilibrium constant for a reaction
 - d. The presence of a catalyst changes the energy of the products and reactants in a reaction
 - e. The presence of a catalyst changes the activation energy for a reaction
- 6. Which of the following is *false* regarding equilibrium?
 - a. Equilibrium can be shifted by changing pressure or temperature
 - b. The forward and reverse reactions stop when a system reaches equilibrium
 - c. The concentrations of products and reactants does not change once the reaction has reached equilibrium

Page 2


- d. The rates of the forward and reverse reactions are equal
- e. Equilibrium concentrations do not depend upon whether you approach equilibrium from the left or the right

- 7. Which of the following is *false* regarding reaction mechanisms?
 - a. Catalysts can appear in the steps of a mechanism
 - b. A mechanism must be composed of elementary reactions
 - c. The steps of the mechanism can contain chemical species that do not appear in the overall correctly balanced chemical equation
 - d. The observed rate law is equal to the sum of the rate laws from all steps
 - e. The observed rate law must agree with the rate law of the slowest step
- 8. For the generic equation

$$aA(g) + bB(g) \iff cC(g) + dD(g)$$

The value of the equilibrium constant, K_c :

- a. Must be measured, it cannot be derived from the balanced equation
- b. Is equal to $([C]^c[D]^d)/([A]^a[B]^b)$
- c. Is equal to $k[A]^a[B]^b$
- d. Is equal to $([A]^a[B]^b)/([C]^c[D]^d)$
- e. Is not affected by temperature



- 9. The reaction represented by the plots above:
 - a. Is zero order
 - b. Is first order
 - c. Is second order
 - d. Is third order
 - e. The order can't be determined by these graphs
- 10. Which of the following statements is *false* regarding the reaction quotient, Q?
 - a. It tells the direction that the reaction must shift to reach equilibrium
 - b. If $Q < K_c$, the system needs to shift toward the reactants to reach equilibrium
 - c. If $Q=K_c$, the system is at equilibrium
 - d. If $\widetilde{Q} > K_c$, the system needs to shift toward the reactants to reach equilibrium
 - e. It has the same mathematical form as the equilibrium constant

Multiple Choice Calculations: (12pts each)

- 11. A reaction is found to be second order with respect to fluoride ion, a reactant. If $[F]_0 =$ 2.91M and $k = 7.47 \times 10^{-3} \text{ M}^{-1} \text{sec}^{-1}$, what will the concentration be after 17 minutes have passed?
 - a. 7.96 M
 - b. 2.12 M
 - c. 0.471 M
 - d. 0.137 M
 - e. 0.126 M
- 12. For the reaction:

 $CH_4(g) + H_2O(g) \iff CO(g) + 3 H_2(g) \qquad \Delta H = 206.2^{kJ}/_{mol}$

The equilibrium concentrations have been found to be $[CO]_{eq} = 1.39M$, $[H_2]_{eq} = 3.27M$, $[CH_4]_{eq} = 4.54 \times 10^{-3} \text{ M}, [H_2O]_{eq} = 8.19 \times 10^{-4} \text{ M}.$ What is the equilibrium constant?

- a. 7.65×10^{-8}
- b. 6.78×10^3
- c. 1.22×10^6
- d. 3.67×10^6
- e. 1.31×10^7
- 13. You have found the following value in a table of equilibrium constants at 25°C:

$$Zn^{2+}(aq) + 4 NH_3(aq) = [Zn(NH_3)_4]^{2+}(aq)$$

$$K_c = 3.6 \times 10^{-10}$$

What is the equilibrium constant for the reaction:

$$3 \left[\text{Zn(NH}_3)_4 \right]^{2+} (\text{aq}) \iff 3 \text{Zn}^{2+} (\text{aq}) + 12 \text{NH}_3 (\text{aq})$$

- a. 2.1×10^{28}
- b. 2.8×10^9
- c. $9.3x10^8$
- d. 1.1x10⁻⁹
- e. 4.7×10^{-29}
- 14. A reaction is found to be first order with respect to reactant A and second order with respect to reactant B. If $[A]_0 = 0.191M$, $[B]_0 = 0.228M$ and $k = 7.38 \times 10^{-2} \,\text{M}^{-2} \text{sec}^{-1}$, what is the initial rate of the reaction?

 - a. $6.14x10^{-4} \text{ M/}_{\text{sec}}$ b. $7.33x10^{-4} \text{ M/}_{\text{sec}}$
 - c. $3.21 \times 10^{-3} \text{ M/}_{\text{sec}}$
 - d. $1.69^{\text{ M}}/_{\text{sec}}$
 - e. $7.43 \, {\rm M/_{sec}}$

Problems: (21pts each)

15. For the reaction:

$$\operatorname{Sn}^{2+}(\operatorname{aq}) + 3 \operatorname{OH}^{-}(\operatorname{aq}) \rightarrow [\operatorname{Sn}(\operatorname{OH})_{3}]^{-}(\operatorname{aq})$$

You have collected the following data at 12.64°C:

Experiment	$[\operatorname{Sn}^{2+}]_{o}$	[OH ⁻] ₀	Rate _{observed}
1	0.481 M	1.143 M	$2.18 \times 10^{-2} \text{ M/}_{\text{sec}}$
2	0.962 M	1.143 M	$8.72 \times 10^{-2} \text{ M/}_{\text{sec}}$
3	0.481 M	3.429 M	$6.54 \times 10^{-2} \text{ M/}_{\text{sec}}$

What are the rate law and the value of the rate law constant, k, for this reaction? If you redo Experiment 2 at 31.56°C, the rate is 0.323 M/sec. What is the activation energy for this reaction?

- 16. When 1.624g of nitrogen dioxide $\{NO_2(g)\}$ and 1.006g of fluorine gas $\{F_2(g)\}$ are sealed together in a 1.500L vessel, they reach equilibrium with nitrogen trifluoride $\{NF_3(g)\}$ and oxygen $\{O_2(g)\}$. If the equilibrium concentration of $F_2(g)$ is found to be 1.686x10⁻⁷ M:
 - a. What are the equilibrium concentrations of all products and reactants?
 - b. What is the value of K_c ?
 - c. Is the reaction Product-favored or reactant-favored?