

Chemistry 210

Exam 1

Be sure to put your name on each page. This page can be removed from your exam so that you will have a Periodic Table handy throughout the exam, it does not need to be turned in. Show all your work for problems which require any sort of calculation, no credit will be given for answers without work shown. If you have shown a significant amount of work or multiple drawings for a problem, draw a box around what you consider your final answer.

Avogadro's Number = 6.022×10^{23} units/mol

$32.00^\circ\text{F} = 0.000^\circ\text{C} = 273.15\text{K}$

Density of Water = $1.000^{\text{g}}/\text{mL}$

$R = 0.08206 \text{ L}\cdot\text{atm}/\text{mol}\cdot\text{K} = 8.314 \text{ J}/\text{mol}\cdot\text{K}$

$PV = nRT$

$\Delta T_{\text{fp/bp}} = k_{\text{fp/bp}} \cdot m \cdot i$

For water: $k_{\text{fp}} = -1.86^\circ\text{C}/m$
 $k_{\text{bp}} = 0.512^\circ\text{C}/m$

$P_1 = X_1 P_1^\circ$

$\Pi = cRTi$

$C_1 V_1 = C_2 V_2$

Quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Integrated Rate Laws:

$\ln[A]_t = -kt + \ln[A]_o$

$1/[A]_t = kt + 1/[A]_o$

$[A]_t = -kt + [A]_o$

$k = Ae^{-E_a/RT}$

$\ln(k) = \left(\frac{-E_a}{R} \right) \left(\frac{1}{T} \right) + \ln(A)$

$\ln\left(\frac{k_1}{k_2} \right) = \frac{E_a}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$

$\text{pH} = \text{pK}_a + \log\left(\frac{[\text{conjugate base}]}{[\text{conjugate acid}]} \right)$

$E_{\text{cell}} = E^\circ_{\text{cell}} - \frac{RT}{nF} \ln Q$

$E^\circ_{\text{cell}} = \frac{RT}{nF} \ln K^\circ$

$K^\circ = e^{(nF/RT) E^\circ_{\text{cell}}}$

$F = 96485 \text{ J}/\text{V}\cdot\text{mol of electrons}$

$\Delta G^\circ = \Delta H^\circ_{\text{system}} - T\Delta S^\circ_{\text{system}}$

$\Delta G^\circ = -nFE^\circ_{\text{cell}} = -RT \ln K^\circ$

$\Delta G = \Delta G^\circ + RT \ln Q$

$F = 96485 \text{ C}/\text{mol electrons}$

$1A = 1 \text{ C} / \text{sec}$

1 H 1.0079																	2 He 4.0026				
3 Li 6.941	4 Be 9.0122															5 B 10.811	6 C 12.011	7 N 14.007	8 O 15.999	9 F 18.998	10 Ne 20.180
11 Na 22.990	12 Mg 24.305															13 Al 26.982	14 Si 28.086	15 P 30.974	16 S 32.066	17 Cl 35.453	18 Ar 39.948
19 K 39.098	20 Ca 40.078	21 Sc 44.956	22 Ti 47.88	23 V 50.942	24 Cr 51.996	25 Mn 54.938	26 Fe 55.847	27 Co 58.933	28 Ni 58.69	29 Cu 63.546	30 Zn 65.39	31 Ga 69.723	32 Ge 72.61	33 As 74.922	34 Se 78.96	35 Br 79.904	36 Kr 83.80				
37 Rb 85.468	38 Sr 87.62	39 Y 88.906	40 Zr 91.224	41 Nb 92.906	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.76	52 Te 127.60	53 I 126.90	54 Xe 131.29				
55 Cs 132.91	56 Ba 137.33	57 La 138.91	72 Hf 178.49	73 Ta 180.95	74 W 183.84	75 Re 186.21	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.97	80 Hg 200.59	81 Tl 204.38	82 Pb 207.2	83 Bi 208.98	84 Po (209)	85 At (210)	86 Rn (222)				
87 Fr (223)	88 Ra 226.03	89 Ac 227.03	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs (265)	109 Mt (266)	110 (269)	111 (272)	112 (277)	114		116							

58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.97	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.94	70 Yb 173.04	71 Lu 174.97
90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np 237.05	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (258)	101 Md (258)	102 No (259)	103 Lr (260)

Multiple Choice (5pts each): Circle the letter of the most correct response.

- Which of the following is **not** a correct ideal gas relationship?
 - $T_1n_1=T_2n_2$
 - $V_1/T_1=V_2/T_2$
 - $T_1/P_1=T_2/P_2$
 - $PV=nRT$
 - $P_1n_1=P_2n_2$**
- Rank the 3 states of matter from lowest kinetic energy to highest kinetic energy.
 - Gas, liquid, solid
 - Solid, liquid, gas**
 - Gas, solid, liquid
 - Liquid, gas, solid
 - Solid, gas, liquid
- Which of the following statements is most correct about colligative properties of an ideal solution?
 - The presence of a solute lowers the vapor pressure of a solution.
 - The presence of a solute raises the boiling point of a solution.
 - The presence of a solute lowers the freezing point of a solution.
 - Colligative properties depend upon the number of solute particles, not on the identity of the solute particles.
 - These statements are all correct.**
- All of the following concentration units require that you use the molar mass of the solute except:
 - Molarity
 - Mass percent**
 - Mole fraction
 - Normality
 - Molality
- When dissolving a solid in a liquid:
 - Formation of solvent-solute interactions is endothermic
 - The boiling point of the solution will be higher than that of the pure solvent**
 - Energy is released (exothermic) by breaking solvent-solvent and solute-solute interactions
 - The enthalpy of solution is always positive
 - The freezing point of the solution will be higher than that of the pure solvent
- The volume of a gas:
 - Increases as the pressure increases
 - Decreases as the kinetic energy increases
 - Is always a constant
 - Increases as the temperature increases**
 - Remains constant as the amount of gas is increased
- Carbon dioxide (CO₂) has a lower boiling point than sulfur dioxide (SO₂) because:
 - The bonds in SO₂ are polar but the bonds in CO₂ are not
 - CO₂ has stronger London dispersion forces than SO₂
 - SO₂ is a polar molecule but CO₂ is not**
 - SO₂ forms stronger hydrogen bonds than CO₂
 - CO₂ sublimates

8. A laboratory technician prepares a solution by weighing out 39.225g of potassium bromide and dissolving it in enough water to make 150.00mL of solution. The technician labels the solution “1.3m KBr(aq)”. Why is this not correct? What should the laboratory technician do to correct the error? Calculate a correct concentration for this solution. (10pts)

The technician diluted the solution to a known volume, so the correct concentration unit (given the information in the problem) can only be molarity, the label should be “M”, not “m”. In addition, the technician only reports 2 sig figs, but the data listed in the problem should permit 5 sig figs.

$$\text{Molarity of KBr(aq)} = \left(\frac{\left(\frac{39.225 \text{g KBr}}{119.002 \frac{\text{g}}{\text{mol}}} \right)}{0.15000 \text{L solution}} \right) = 2.1974 \text{M KBr(aq)}$$

It looks like not only was the unit wrong, but the numerical value was wrong as well, it appears that the technician used the formula “KBr₂” when calculating the formula weight. That technician should look for a different job...

9. You have prepared a solution by dissolving 21.918g of potassium phosphate in enough water to make 500.0mL of solution. What is the *molarity* of this solution? (10pts)

$$\text{Molarity of K}_3\text{PO}_4(\text{aq}) = \left(\frac{\left(\frac{21.918 \text{g K}_3\text{PO}_4}{212.264 \frac{\text{g}}{\text{mol}}} \right)}{0.5000 \text{L solution}} \right) = 0.2065 \text{M K}_3\text{PO}_4(\text{aq})$$

10. You have prepared a solution by dissolving 9.986g of ammonium iodide in 100.0g of water. What is the *molality* of this solution? (10pts)

$$\text{Molality of NH}_4\text{I(aq)} = \left(\frac{\left(\frac{9.986 \text{g NH}_4\text{I}}{144.94 \frac{\text{g}}{\text{mol}}} \right)}{0.1000 \text{kg solvent}} \right) = 0.6890 \text{m NH}_4\text{I(aq)}$$

11. You have prepared a solution by diluting 50.00mL of a 2.312M aqueous solution of sugar (C₆H₁₂O₆) to a total volume of 200.0mL. What is the *molarity* of this solution? (10pts)

For dilutions, we can use the formula $C_1V_1 = C_2V_2$.

$$(2.312 \text{M})(50.00 \text{mL}) = C_2(200.0 \text{mL})$$

$$C_2 = 0.5780 \text{M}$$

12. A 2.00L cylinder contains helium gas at 18.93°C and 2.64atm pressure. How many grams of He are in the cylinder? (10pts)

This is a single-condition ideal gas law problem, so we should be able to solve it using $PV=nRT$.

$$PV = nRT$$
$$(2.64\text{atm})(2.00\text{L}) = n(0.08206 \text{ L}\cdot\text{atm}/\text{mol}\cdot\text{K})(292.08\text{K})$$
$$n = 0.22029\text{mols (I haven't rounded to correct sig figs yet...)}$$
$$(0.22029\text{mols})(4.0026 \text{ g/mol}) = 0.882\text{g He}$$

13. What is the freezing point of a solution made by dissolving 22.734g of lithium nitrate in 200.0g of water? (15pts)

$$\text{Molality of LiNO}_3(\text{aq}) = \left(\frac{\left(\frac{22.734\text{g LiNO}_3}{68.945 \text{ g/mol}} \right)}{0.2000\text{kg solvent}} \right) = 1.6487\text{m LiNO}_3(\text{aq})$$

$$\Delta T_{\text{fp}} = k_{\text{fpd}} \cdot m \cdot n$$
$$\Delta T_{\text{fp}} = (1.86 \text{ }^\circ\text{C}/\text{m})(1.6487\text{m})(2 \text{ mols particles/mol LiNO}_3)$$
$$\Delta T_{\text{fp}} = 6.13^\circ\text{C}$$

Since freezing point is depressed, the solution will freeze 6.13°C lower than the freezing point of the pure solvent, in this case -6.13°C .

14. Each of the following solids is dissolved in separate beakers containing 500.0mL of water. Rank the solutions from lowest vapor pressure to highest vapor pressure and explain your answer. (15pts)
- 0.4mols magnesium phosphate
 - 0.6mols sodium chloride
 - 0.5mols calcium nitrate
 - 0.7mols ammonium phosphate

Vapor pressure depression is a colligative property, so the amount that the P_{vap} is depressed depends upon the number of solute particles, not their identity. Let's look at how many solute particles there will be for each of the listed solutes..

0.4mols of $\text{Mg}_3(\text{PO}_4)_2$ dissolve to give $3(0.4\text{mol})=1.2\text{mol}$ of $\text{Mg}^{2+}(\text{aq})$ ions and $2(0.4\text{mol})=0.8\text{mol}$ of $\text{PO}_4^{3-}(\text{aq})$ ions, for a total of $1.2+0.8=2.0\text{mols}$ of solute particles

0.6mols of NaCl yields 0.6mol of $\text{Na}^+(\text{aq})$ and 0.6mol of $\text{Cl}^-(\text{aq})$, so 1.2mols of solute particles

0.7mols of $(\text{NH}_4)_3\text{PO}_4$ yields $(3 \times 0.7)+0.7=2.8\text{mols}$ of solute particles

0.5mols of $\text{Ca}(\text{NO}_3)_2$ yields $3 \times 0.5=1.5\text{mols}$ of solute particles

The more solute particles, the lower the vapor pressure of the solution, so the ranking should be:

$$0.7\text{mols of } (\text{NH}_4)_3\text{PO}_4 < 0.4\text{mols of } \text{Mg}_3(\text{PO}_4)_2 < 0.5\text{mols of } \text{Ca}(\text{NO}_3)_2 < 0.6\text{mols of } \text{NaCl}$$

15. You have a 63.926g sample of steam (gaseous water) at 142.19°C. Describe what happens to this sample when it is cooled to 38.63°C and calculate the amount of energy transferred during cooling. (15pts)

$$\{C_s(\text{ice}) = 2.09 \text{ J/g}\cdot\text{K}; C_s(\text{water}) = 4.184 \text{ J/g}\cdot\text{K}; C_s(\text{steam}) = 2.01 \text{ J/g}\cdot\text{K}; \Delta H_{\text{fusion}}(\text{water}) = 6.02 \text{ kJ/mol}; \Delta H_{\text{vaporization}}(\text{water}) = 40.7 \text{ kJ/mol}\}$$

From 142.19°C to 100°C, the steam is cooling. At 100°C, the steam condenses. From 100°C to 38.63°C, the liquid water is cooling.

Heat capacity of steam:	$(2.01 \text{ J/g}\cdot\text{K})(63.926\text{g})(42.19\text{K})(\frac{1\text{kJ}}{1000\text{J}}) = 5.42\text{kJ}$
Phase change:	$(40.7 \text{ kJ/mol})(63.926\text{g}) / (18.015 \text{ g/mol}) = 144\text{kJ}$
Heat capacity of water:	$(4.184 \text{ J/g}\cdot\text{K})(63.926\text{g})(61.37\text{K})(\frac{1\text{kJ}}{1000\text{J}}) = 16.41\text{kJ}$
Total energy transferred:	166kJ

16. A newly discovered protein has been isolated from seeds of a tropical plant and needs to be characterized. A total of 0.137g of this protein was dissolved in enough water to produce 2.00mL of solution. At 31.68°C the osmotic pressure produced by the solution was 0.134atm. What is the molar mass of the protein? (20pts)

This is actually one of the most common ways that osmotic pressure changes are used in real laboratory settings...

$$\Pi = MRT$$

$$(0.134\text{atm}) = \left(\frac{\left(\frac{0.137\text{g}}{x \text{ g/mol}} \right)}{0.00200\text{L solution}} \right) (0.08206 \text{ L}\cdot\text{atm/mol}\cdot\text{K})(304.83\text{K})$$

$$x = 12800 \text{ g/mol}$$

We're assuming that this protein does not dissociate in solution, so $n=1$. Proteins are very large molecules, so this molar mass is reasonable.