Chemistry 210

Exam 2

Be sure to put your name on each page. This page can be removed from your exam so that you will have a Periodic Table handy throughout the exam, it does not need to be turned in. Show all your work for problems which require any sort of calculation, no credit will be given for answers without work shown. If you have shown a significant amount of work or multiple drawings for a problem, draw a box around what you consider your final answer.

Avogadro's Number = 6.022×10^{23} units/mol $32.00^{\circ}F = 0.000^{\circ}C = 273.15K$ Density of Water = $1.000^{g}/_{mL}$ R = 0.08206 L*atm/mol*K = 8.314 J/mol*K = 8.3

Integrated Rate Laws:
$$0^{th} \text{ order } [A]_t = -kt + [A]_o$$

$$1^{st} \text{ order } \ln[A]_t = -kt + \ln[A]_o$$

$$2^{nd} \text{ order } 1/[A]_t = kt + 1/[A]_o$$

$$k = Ae^{-Ea/RT}$$

$$\ln(k) = \left(\frac{-E_a}{R}\right) \left(\frac{1}{T}\right) + \ln(A)$$

$$\ln\left(\frac{k_1}{k_2}\right) = \frac{E_a}{R} \left(\frac{1}{T_2} - \frac{1}{T_1}\right)$$

$$pH = pK_a + \log\left(\frac{[\text{conjugate base}]}{[\text{conjugate acid}]}\right)$$

$E_{cell} = E^{o}_{cell} - {^{RT}}/_{nF} \ln Q$
$E_{cell}^{o} = {^{RT}}/{_{nF}} lnK^{o}$
$K^{o} = e^{(nF)}_{RT} E^{o}_{cell}$
$F = 96485 ^{J}/_{V \bullet mol \ of \ electrons}$
$\Delta G^{o} = \Delta H^{o}_{system} - T\Delta S^{o}_{system}$
$\Delta G^{o} = -nFE^{o}_{cell} = -RTlnK^{o}$
$\Delta G = \Delta G^{o} + RT lnQ$
$F = 96485$ $^{\rm C}/_{\rm mol\ electrons}$
1A = 1 C / sec

1																	2
H																	He
1.0079	4	1										5	6	7	8	9	4.0026
Li	Be											\mathbf{B}	$\overset{\circ}{\mathbf{C}}$	Ń	o	F	Ne
6.941	9.0122											10.811	12.011	14.007	15.999	₽ 18.998	20.180
11	12											13	14	15	16	17	18
Na	Mg											Al	Si	P	S	Cl	Ar
22.990	24.305											26.982	28.086	■ 30.974	32.066	35.453	39.948
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K	Ca	Sc	Ti	\mathbf{V}	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
39.098	40.078	44.956	47.88	50.942	51.996	54.938	55.847	58.933	58.69	63.546	65.39	69.723	72.61	74.922	78.96	79.904	83.80
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
85.468	87.62	88.906	91.224	92.906	95.94	(98)	101.07	102.91	106.42	107.87	112.41	114.82	118.71	121.76	127.60	126.90	131.29
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ba	La	Hf	Ta	\mathbf{W}	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
132.91	137.33	138.91	178.49	180.95	183.84	186.21	190.23	192.22	195.08	196.97	200.59	204.38	207.2	208.98	(209)	(210)	(222)
87	88	89	104	105	106	107	108	109	110	111	112		114		116		•
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt									
(223)	226.03	227.03	(261)	(262)	(263)	(262)	(265)	(266)	(269)	(272)	(277)						

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
140.12	140.91	144.24	(145)	150.36	151.97	157.25	158.93	162.50	164.93	167.26	168.94	173.04	174.97
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	\mathbf{U}	Np	Pu	Am	Cm	$\mathbf{B}\mathbf{k}$	Cf	Es	Fm	Md	No	Lr
232.04	231.04	238.03	237.05	(244)	(243)	(247)	(247)	(251)	(252)	(258)	(258)	(259)	(260)

Page 1


Multiple Choice: Circle the letter of the most correct response. (5pts per question)

- 1. Which of the following does *not* affect the rate of a reaction?
 - a. The coefficients of the reactants in the balanced equation
 - b. The temperature of the system
 - c. The energy of collisions between reacting particles
 - d. The frequency of collisions between reacting particles
 - e. The orientation of colliding particles
- 2. For the generic equation:

$$aA(g) + bB(g) \rightarrow cC(g) + dD(g)$$

Which of the following is a correct expression of the rate of the reaction:

- a. $\frac{1}{a} \frac{\Delta[A]}{\Delta}$
- b. $k[A]^a[B]^b$
- c. $^{-1}/_{d} ^{\Delta[D]}/_{\Delta t}$
- d. $k[C]^{c}[D]^{d}$
- e. $\frac{-1}{b} \frac{\Delta[B]}{\Delta[A]}$
- 3. If the rate of a reaction increases by a factor of 4 when the initial concentration of reactant "A" is increased by a factor of 2, the reaction must be:
 - a. Oth order with respect to [A]_o
 - b. 1st order with respect to [A]_o
 - c. 2nd order overall
 - d. 2nd order with respect to [A]_o
 - e. The order of the reaction depends on the balanced chemical equation
- 4. For a first order reaction:
 - a. The slope of the integrated rate law plot is equal to k
 - b. The slope of the integrated rate law plot is equal to $(-E_a/R)$
 - c. The intercept of the integrated rate law is equal to the ln of the initial concentration
 - d. The intercept of the integrated rate law plot is equal to the initial concentration
 - e. The slope of the integrated rate law is equal to the frequency factor, A.



- 5. The reaction represented by the plots above:
 - a. Is zero order
 - b. Is first order
 - c. Is second order
 - d. Is third order
 - e. The order can't be determined by these graphs

- 6. Which of the following is *false* regarding reaction mechanisms?
 - a. The observed rate law is equal to the sum of the rate laws from all steps
 - b. The observed rate law must agree with the rate law of the slowest step
 - c. The steps of the mechanism can contain chemical species that do not appear in the overall correctly balanced chemical equation
 - d. Catalysts can appear in the steps of a mechanism
 - e. A mechanism must be composed of elementary reactions
- 7. Which of the following is *true* regarding catalysts and catalyzed reactions?
 - a. The presence of a catalyst does not change the mechanism of a reaction
 - b. The presence of a catalyst changes the equilibrium constant for a reaction
 - c. The presence of a catalyst changes the activation energy for a reaction
 - d. The presence of a catalyst changes the energy of the products and reactants in a reaction
 - e. The concentration of a catalyst cannot appear in the rate law for a reaction
- 8. Which of the following is *false* regarding equilibrium?
 - a. The concentrations of products and reactants does not change once the reaction has reached equilibrium
 - b. Equilibrium can often be shifted by changing pressure or temperature
 - c. The rates of the forward and reverse reactions are equal
 - d. Equilibrium concentrations do not depend upon whether you approach equilibrium from the left or the right
 - e. The forward and reverse reactions stop when a system reaches equilibrium
- 9. For the generic equation

$$aA(g) + bB(g) = cC(g) + dD(g)$$

The value of the equilibrium constant, K_c:

- a. Is not affected by temperature
- b. Is equal to $([A]^{a}[B]^{b})/([C]^{c}[D]^{d})$
- c. Is equal to k[A]^a[B]^b
- d. Is equal to $([C]^c[D]^d)/([A]^a[B]^b)$
- e. Must be measured, it cannot be derived from the balanced equation
- 10. Which of the following is *true* regarding equilibrium reactions?
 - a. If K < 0, the reaction reaches equilibrium very quickly.
 - b. If K > 1, the reaction is reactant-favored.
 - c. If K = 1, the reaction has stopped.
 - d. If K > 1, the reaction is product-favored.
 - e. If K is very small, the limiting reactant is very nearly used up.
- 11. Considering the reaction given, all of the following stresses will shift the equilibrium to the right except:

$$CO(g) + H_2O(g) \leftrightarrow CO_2(g) + H_2(g)$$
 $\Delta H_{rxn} = 131 \text{ }^{kJ}/_{mol}$

- a. Removing carbon dioxide from the system
- b. Increasing the temperature of the system
- c. Adding carbon monoxide to the system
- d. Increasing the pressure on the system
- e. Removing hydrogen from the system
- 12. Which of the following statements is *true* regarding the reaction quotient, Q?
 - a. It indicates how quickly a reaction will reach equilibrium
 - b. If $Q > K_c$, the system needs to shift toward the reactants to reach equilibrium
 - c. If $Q=K_c$, the reaction has stopped
 - d. If Q>K_c, the system needs to shift toward the products to reach equilibrium
 - e. It has the same mathematical form as the rate law expression

Problems: Show your work.

13. A reaction is found to be second order with respect to reactant A and zero order with respect to reactant B. If $[A]_o = 0.351M$, $[B]_o = 0.442M$ and $k = 5.18 \times 10^{-3} M^{-1} sec^{-1}$, what is the initial rate of the reaction? (12pts)

14. For the reaction:

 $2 \text{ CH}_3\text{OH}(g) + 3 \text{ F}_2(g) \iff 2 \text{ CF}_3\text{OH}(g) + 3 \text{ H}_2(g) \qquad \Delta H = -437.2 \text{ kJ}/_{mol}$ The equilibrium concentrations are observed: $[\text{CH}_3\text{OH}]_{eq} = 0.291\text{M}$, $[\text{F}_2]_{eq} = 0.118\text{M}$, $[\text{CF}_3\text{OH}]_{eq} = 7.72 \times 10^{-2} \text{ M}$, $[\text{H}_2]_{eq} = 0.834 \text{ M}$. What is the equilibrium constant for this reaction? (12pts)

15. A reaction is found to be zero order with respect to carbonate ion, a reactant. If $[CO_3^{-2}]_0 = 3.94M$ and $k = 6.81 \times 10^{-2}$ M•min⁻¹, how much time must pass before the concentration of carbonate ions falls to $1.54 \times 10^{-3} M$? (12pts)

16. You have found the following value in a table of equilibrium constants at 25°C:

$$Fe^{3+}(aq) + 4 Cl^{-}(aq) = [FeCl_4]^{-1}(aq)$$
 $K_c = 2.16x10^5$

What is the equilibrium constant for the following reaction? (12pts)

$$2 [FeCl4]-1(aq) \iff 2 Fe3+(aq) + 8 Cl-(aq)$$

17. For the reaction:

$$2 \text{ NO}_2(g) + \text{Br}_2(aq) \rightarrow 2 \text{ NOBr}(g) + \text{O}_2(g)$$

You have collected the following data at 46.92°C:

Experiment	$[NO_2]_o$	$[Br_2]_{o}$	Rate _{observed}
1	1.38 M	1.62 M	$1.37 \times 10^{-2} \text{ M/}_{min}$
2	2.76 M	1.62 M	$1.37 \times 10^{-2} \text{ M/}_{min}$
3	1.38 M	3.24 M	$5.48 \times 10^{-2} \text{ M/}_{min}$

What are the rate law and the value of the rate law constant, k, for this reaction? If you redo Experiment 3 at 16.31° C, the rate is 7.53×10^{-4} M/_{min}. What is the activation energy for this reaction? (21pts)

- 18. When 0.267mols of sulfur dioxide $\{SO_2(g)\}$ and 0.338mols of fluorine gas $\{F_2(g)\}$ are sealed together in a 1.500L vessel, they reach equilibrium with thionyl fluoride $\{SOF_2(g)\}$ and oxygen $\{O_2(g)\}$. The equilibrium concentration of $F_2(g)$ is found to be 0.193 M. (21pts)
 - a. What are the equilibrium concentrations of all products and reactants?
 - b. What is the value of K_c ?
 - c. Is the reaction product-favored or reactant-favored?