Chemistry 210

Exam 3

Be sure to put your name on each page. This page can be removed from your exam so that you will have a Periodic Table handy throughout the exam, it does not need to be turned in. Show all your work for problems which require any sort of calculation, no credit will be given for answers without work shown. If you have shown a significant amount of work or multiple drawings for a problem, draw a box around what you consider your final answer.

Avogadro's Number =
$$6.022 \times 10^{23}$$
 units/ $_{mol}$ 32.00°F = 0.000 °C = 273.15 K Density of Water = $1.000^g/_{mL}$ R = 0.08206 L*atm/ $_{mol*K}$ = 8.314 J/ $_{mol*K}$ PV=nRT $\Delta T_{fp/bp} = k_{fp/bp}$ *m*i For water: $k_{fp} = -1.86$ °C/ $_{m}$ $k_{bp} = 0.512$ °C/ $_{m}$ $\Pi = MRTi$ $C_1V_1 = C_2V_2$

Quadratic formula:

 $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

$$\begin{split} & In[A]_t = -kt + In[A]_o \\ & 1/[A]_t = kt + 1/[A]_o \\ & [A]_t = kt + 1/[A]_o \\ & [A]_t = -kt + [A]_o \\ & k = Ae^{-Ea/RT} \\ & ln(k) = \left(\frac{-E_a}{R}\right) \left(\frac{1}{T}\right) + ln(A) \\ & ln\left(\frac{k_1}{k_2}\right) = \frac{E_a}{R} \left(\frac{1}{T_2} - \frac{1}{T_1}\right) \\ & pH = pK_a + log\left(\frac{[conjugate\ base]}{[conjugate\ acid]}\right) \end{split}$$

$$\begin{split} E_{cell} &= E^{o}_{cell} - {^{RT}}\!/_{nF} \, lnQ \\ E^{o}_{cell} &= {^{RT}}\!/_{nF} \, lnK^{o} \\ K^{o} &= e^{\wedge}({^{nF}}\!/_{RT} \, E^{o}_{cell}) \\ F &= 96485 \, {^{J}}\!/_{v \cdot mol \ of \ electrons} \\ \Delta G^{o} &= \Delta H^{o}_{system} - T\Delta S^{o}_{system} \\ \Delta G^{o} &= -nFE^{o}_{cell} = -RT lnK^{o} \\ \Delta G &= \Delta G^{o} + RT lnQ \\ F &= 96485 \, {^{C}}\!/_{mol \ electrons} \\ 1A &= 1 \, C \, / \, sec \end{split}$$

	1																
1																	2
H																	He
1.0079																	4.0026
3	4											5	6	7	8	9	10
Li	Be											В	C	N	O	\mathbf{F}	Ne
6.941	9.0122											10.811	12.011	14.007	15.999	18.998	20.180
11	12											13	14	15	16	17	18
Na	Mg											Al	Si	P	S	Cl	Ar
22.990	24.305											26.982	28.086	30.974	32.066	35.453	39.948
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K	Ca	Sc	Ti	\mathbf{V}	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
39.098	40.078	44.956	47.88	50.942	51.996	54.938	55.847	58.933	58.69	63.546	65.39	69.723	72.61	74.922	78.96	79.904	83.80
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb	Sr	${f Y}$	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	Ι	Xe
85.468	87.62	88.906	91.224	92.906	95.94	(98)	101.07	102.91	106.42	107.87	112.41	114.82	118.71	121.76	127.60	126.90	131.29
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ba	La	Hf	Ta	\mathbf{W}	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
132.91	137.33	138.91	178.49	180.95	183.84	186.21	190.23	192.22	195.08	196.97	200.59	204.38	207.2	208.98	(209)	(210)	(222)
87	88	89	104	105	106	107	108	109	110	111	112		114		116		
Fr	Ra	\mathbf{Ac}	Rf	Db	Sg	Bh	Hs	Mt									
(223)	226.03	227.03	(261)	(262)	(263)	(262)	(265)	(266)	(269)	(272)	(277)						

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
140.12	140.91	144.24	(145)	150.36	151.97	157.25	158.93	162.50	164.93	167.26	168.94	173.04	174.97
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	\mathbf{U}	Np	Pu	Am	Cm	$\mathbf{B}\mathbf{k}$	Cf	Es	Fm	Md	No	Lr
232.04	231.04	238.03	237.05	(244)	(243)	(247)	(247)	(251)	(252)	(258)	(258)	(259)	(260)

1. Complete each row of the following tables for aqueous solutions at 25°C (4pts per box):

$[H_3O^+]$	[OH ⁻]	рН	рОН	Acidic, Basic or Neutral?	
	3.82x10 ⁻¹¹				
		11.158			

Conjugate Acid	K _a @25°C	Conjugate Base	К _b @25°С
H ₂ SeO ₃			$4.3x10^{-12}$
	1.0x10 ⁻⁷	HAsO ₄ ² -	

2. Does the combination listed result in an effective buffer solution? (4pts each)

Yes No 0.64mol HClO₄(aq) + 0.68mol KOH(aq)

Yes No 0.90mol Na₂HPO₄(aq) + 0.45mol HNO₃(aq)

Yes No 1.28mol $Na_2CO_3(aq) + 1.92mol HCl(aq)$

Yes No 0.93mol CH₃COOH(aq) + 1.06mol CH₃COOK(aq)

Yes No 0.06mol $HNO_2(aq) + 1.18$ mol $LiNO_2(aq)$

3. How much 0.583M KOH(aq) must be added to 25.00mL of 0.332M HBr(aq) to reach the equivalence point? What is the pH of this solution at the equivalence point? Explain. (10pts)

4. You have prepared a buffer solution by combining 0.482mols of chlorous acid (HClO₂, $K_a = 1.2 \times 10^{-2}$) and 0.513mols of potassium chlorite in enough water to make 500.0mL of solution. What is the pH of this buffer solution? (15pts)

5. What is the K_a of a weak acid if 500.0mL of a solution containing 0.278mol of the acid and 0.327 mol of its conjugate base has a pH of 8.243? Over what pH range would this conjugate acid/ conjugate base pair make an effective buffer? (15pts)

- 6. You are going to use a titration to determine the concentration of an unknown potassium sulfite solution (K_2SO_3 , $pK_b = 6.806$). The acid you have chosen to use is 0.927M perchloric acid.
 - a. Write out the chemical equations for the step-wise protonation/neutralization of sulfite and the overall/net chemical reaction. (6pts)

b. Someone left your pH probe laying on the bench and it has dried out. After a short search, you have found the following three visual indicators: Thymolphthalein (TP, endpoint pH range = 9.5-10.5); Phenol red (PR, endpoint pH range = 6.8-8.3); Bromophenol Blue (BPB, endpoint pH range = 3.1-4.6). Which of these visual indicators would be useful in your titration? Explain your choice(s). (12pts)

Page 3

(#6 continued)

c. Sketch the titration curve you would expect for this titration, labeling all equivalence points and sulfite-based species present in each portion of the curve. (6pts)

d. As you are about to start your titration, a package arrives with a new pH probe, so you decide to use the pH probe to monitor your titration. You find that 25.00mL of potassium sulfite requires 34.62mL of perchloric acid to reach the first equivalence point. What is the concentration of the potassium sulfite solution? (12pts)

e. How much perchloric acid solution would be required to reach the second equivalence point in part "d"? Explain your answer. (6pts)