Chemistry 210

Exam 2

Be sure to put your name on each page. This page can be removed from your exam so that you will have a Periodic Table handy throughout the exam, it does not need to be turned in. Show all your work for problems which require any sort of calculation, no credit will be given for answers without work shown. If you have shown a significant amount of work or multiple drawings for a problem, draw a box around what you consider your final answer.

$$\begin{array}{l} A vogadro's \ Number = 6.022x10^{23 \ units}/_{mol} \\ 32.00°F = 0.000°C = 273.15K \\ Density \ of \ Water = 1.000^g/_{mL} \\ R = 0.08206^{\ L*atm}/_{mol*K} = 8.314^{\ J}/_{mol*K} \\ PV = nRT \\ \Delta T_{fp/bp} = k_{fp/bp} \bullet m \bullet i \\ For \ water, \ k_{fp} = -1.86^{°C}/_{m} \ ; \ k_{bp} = 0.52^{°C}/_{m} \\ P_1 = X_1 P_1° \\ P = cRTi \\ C_1 V_1 = C_2 V_2 \\ Quadratic \ formula: \\ x = \frac{-b \pm \sqrt{b^2 - 4ac}}{1} \\ \end{array}$$

$$\begin{split} & \text{Integrated Rate Laws:} \\ & 0^{\text{th}} \text{ order} \quad [A]_t = -kt + [A]_o \\ & 1^{\text{st}} \text{ order} \quad \ln[A]_t = -kt + \ln[A]_o \\ & 2^{\text{nd}} \text{ order} \quad 1/[A]_t = kt + 1/[A]_o \\ & k = Ae^{\text{-Ea/RT}} \\ & \ln(k) = \left(\frac{-E_a}{R}\right) \left(\frac{1}{T}\right) + \ln(A) \\ & \ln\left(\frac{k_1}{k_2}\right) = \frac{E_a}{R} \left(\frac{1}{T_2} - \frac{1}{T_1}\right) \\ & pH = pK_a + \log\left(\frac{[\text{conjugate base}]}{[\text{conjugate acid}]}\right) \end{split}$$

$$\begin{split} E_{cell} &= E^{\circ}_{cell} - {^{RT}}/_{nF} \, lnQ \\ E^{\circ}_{cell} &= {^{RT}}/_{nF} \, lnK^{\circ} \\ K^{\circ} &= e^{\wedge}({^{nF}}/_{RT} \, E^{\circ}_{cell}) \\ F &= 96485 \, {^{J}}/_{V^{\bullet}mol \, of \, electrons} \\ \Delta G^{\circ} &= \Delta H^{\circ}_{system} - T\Delta S^{\circ}_{system} \\ \Delta G^{\circ} &= -nFE^{\circ}_{cell} = -RT lnK^{\circ} \\ \Delta G &= \Delta G^{\circ} + RT lnQ \\ F &= 96485 \, {^{C}}/_{mol \, electrons} \\ 1A &= 1 \, C \, / \, sec \end{split}$$

1	1																2
H																	He
1.0079																	4.0026
3	4											5	6	7	8	9	10
Li	Be											В	C	N	O	\mathbf{F}	Ne
6.941	9.0122											10.811	12.011	14.007	15.999	18.998	20.180
11	12											13	14	15	16	17	18
Na	Mg											Al	Si	P	S	Cl	Ar
22.990	24.305											26.982	28.086	30.974	32.066	35.453	39.948
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K	Ca	Sc	Ti	\mathbf{V}	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
39.098	40.078	44.956	47.88	50.942	51.996	54.938	55.847	58.933	58.69	63.546	65.39	69.723	72.61	74.922	78.96	79.904	83.80
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb	Sr	\mathbf{Y}	Zr	Nb	Mo	Tc	Ru	Rh	Pd	$\mathbf{A}\mathbf{g}$	Cd	In	Sn	Sb	Te	I	Xe
85.468	87.62	88.906	91.224	92.906	95.94	(98)	101.07	102.91	106.42	107.87	112.41	114.82	118.71	121.76	127.60	126.90	131.29
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ba	La	Hf	Ta	\mathbf{W}	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
132.91	137.33	138.91	178.49	180.95	183.84	186.21	190.23	192.22	195.08	196.97	200.59	204.38	207.2	208.98	(209)	(210)	(222)
87	88	89	104	105	106	107	108	109	110	111	112		114		116		
0,	00	0,															
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt									

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
140.12	140.91	144.24	(145)	150.36	151.97	157.25	158.93	162.50	164.93	167.26	168.94	173.04	174.97
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
232.04	231.04	238.03	237.05	(244)	(243)	(247)	(247)	(251)	(252)	(258)	(258)	(259)	(260)

Summer 2010

Multiple Choice (7pts each): Circle the letter of the most correct response.

- 1. For a reaction at equilibrium:
 - a. The reactants and products must be in the gas phase.
 - b. The concentration of reactants is equal to the concentration of products.
 - c. The reaction has stopped.
 - d. The mass of reactants is equal to the mass of products.
 - e. The rate of the forward reaction is equal to the rate of the reverse reaction.
- 2. Which of the following is *true* regarding equilibrium reactions?
 - a. If K = 1, the reaction has stopped.
 - b. If K < 0, the reaction reaches equilibrium very quickly.
 - c. If K > 1, the reaction is reactant-favored.
 - d. If K < 1, the reaction is product-favored.
 - e. If K is very large, the limiting reactant is very nearly used up.
- 3. Which of the following is *false* regarding equilibrium?
 - a. Equilibrium can usually be shifted by changing pressure or temperature
 - b. The rates of the forward and reverse reactions are equal
 - c. Equilibrium concentrations do not depend upon whether you approach equilibrium from the left or the right
 - d. The forward and reverse reactions stop when a system reaches equilibrium
 - e. The concentrations of products and reactants does not change once the reaction has reached equilibrium
- 4. Which of the following equilibrium constant expressions is correct for the given reaction:

$$2\;HNO_3(g)\;+\;NO(g)\;\;\leftrightarrow\;\;2\;NO_2(g)\;+\;H_2O(g)$$

a.
$$K_c = \frac{2[NO_2]_{eq}[H_2O]_{eq}}{2[HNO_3]_{eq}[NO]_{eq}}$$

$$b. \quad K_{c} = \frac{\left[NO_{2}\right]_{eq}^{2} + \left[H_{2}O\right]_{eq}}{\left[HNO_{3}\right]_{eq}^{2} + \left[NO\right]_{eq}}$$

$$c. \quad K_c = \frac{\left[NO_2\right]_{eq}^2 \left[H_2O\right]_{eq}}{\left[HNO_3\right]_{eq}^2 \left[NO\right]_{eq}}$$

$$\label{eq:Kc} d. \quad K_c = \frac{\left[HNO_3\right]_{eq}^2\left[NO\right]_{eq}}{\left[NO_2\right]_{eq}^2\left[H_2O\right]_{eq}}$$

$$e. \quad K_c = \frac{\left[NO_2\right]_{eq}^2}{\left[HNO_3\right]_{eq}^2\left[NO\right]_{eq}}$$

- 5. All of the following can be explained by LeChatelier's Principle except:
 - a. Removing a gaseous product will shift the equilibrium right.
 - b. Adding more of an aqueous reactant will shift the equilibrium right.
 - c. Increasing the temperature of an endothermic reaction will shift the equilibrium right.
 - d. Increasing the pressure will shift an equilibrium toward the side that has more gas particles.
 - e. Removing a gaseous reactant will shift the equilibrium left.

Summer 2010

6. Considering the reaction given, all of the following stresses will shift the equilibrium to the left except:

 $CO(g) + H_2O(g) \leftrightarrow CO_2(g) + H_2(g) \qquad \Delta H_{rxn} = 131 \text{ }^{kJ}/_{mol}$

- a. Decreasing the pressure on the system
- b. Adding carbon dioxide to the system
- c. Adding hydrogen to the system
- d. Decreasing the temperature of the system
- e. Removing carbon monoxide from the system
- 7. The reaction quotient for a reaction:
 - a. Tells you how fast the reaction happens
 - b. Is usually a negative number
 - c. Is a constant
 - d. Tells you what direction the reaction must shift to reach equilibrium
 - e. Is the concentration of reactants divided by the concentration of products
- 8. Which of the following statements is *false* regarding the reaction quotient, Q?
 - a. It tells the direction that the reaction must shift to reach equilibrium
 - b. If $Q < K_c$, the system needs to shift toward the products to reach equilibrium
 - c. If $Q=K_c$, the system is at equilibrium
 - d. If Q>K_c, the system needs to shift toward the products to reach equilibrium
 - e. It has the same mathematical form as the equilibrium constant

Problems:

9. You have been studying the reaction of $N_2O(g)$ with $O_2(g)$ to form $NO_2(g)$. After your system has reached equilibrium, you find that the concentrations of all species are: $[N_2O]_{eq} = 0.142M$, $[O_2]_{eq} = 0.216M$, $[NO_2]_{eq} = 1.85M$. What is the value of the equilibrium constant for this reaction? (12pts)

10. A saturated potassium bromide solution is prepared by dissolving KBr in pure water and has a potassium ion concentration of 4.137M at some temperature. What is the K_{sp} of potassium bromide at this temperature? (16pts)

Page 3

11. What are the concentrations of strontium ions (Atomic # = 38) and hydroxide ions in a saturated solution of strontium hydroxide prepared from strontium hydroxide solid in water ($K_{sp} = 3.84 \times 10^{-3}$)? (16pts)

12. NH₃(g) reacts with ClO(g) to form NO₃(g) and HCl(g) with an equilibrium constant of 7.16x10⁻⁷. In a 2.00L reaction vessel, you have combined 0.372mols of NH₃ with 0.546mols of ClO. What are the concentrations of all reactants and products when this reaction reaches equilibrium? (25pts)

- 13. When 0.267mols of sulfur dioxide $\{SO_2(g)\}$ and 0.338mols of fluorine gas $\{F_2(g)\}$ are sealed together in a 1.500L vessel, they reach equilibrium with thionyl fluoride $\{SOF_2(g)\}$ and oxygen $\{O_2(g)\}$. The equilibrium concentration of $F_2(g)$ is found to be 0.193 M. (25pts)
 - a. What are the equilibrium concentrations of all products and reactants?
 - b. What is the value of K_c ?
 - c. Is the reaction product-favored or reactant-favored?