

Chemistry 210

Exam 1

Be sure to put your name on each page. This page can be removed from your exam so that you will have a Periodic Table handy throughout the exam, it does not need to be turned in. Show all your work for problems which require any sort of calculation, no credit will be given for answers without work shown. If you have shown a significant amount of work or multiple drawings for a problem, draw a box around what you consider your final answer.

Avogadro's Number = 6.022×10^{23} units/mol

$32.00^\circ\text{F} = 0.000^\circ\text{C} = 273.15\text{K}$

Density of Water = $1.000^{\text{g}}/\text{mL}$

$R = 0.08206 \text{ L}\cdot\text{atm}/\text{mol}\cdot\text{K} = 8.314 \text{ J}/\text{mol}\cdot\text{K}$

$1\text{atm} = 760\text{torr} = 760\text{mmHg} = 101.325\text{kPa}$

$PV = nRT$

$\Delta T_{\text{fp/bp}} = k_{\text{fp/bp}} \cdot m \cdot i$

For water: $k_{\text{fp}} = -1.86^\circ\text{C}/m$
 $k_{\text{bp}} = 0.512^\circ\text{C}/m$

$P_1 = X_1 P_1^\circ$

$\Pi = MRTi$

$C_1 V_1 = C_2 V_2$

Integrated Rate Laws:

$\ln[A]_t = -kt + \ln[A]_o$

$1/[A]_t = kt + 1/[A]_o$

$[A]_t = -kt + [A]_o$

$k = Ae^{-E_a/RT}$

$\ln(k) = \left(\frac{-E_a}{R} \right) \left(\frac{1}{T} \right) + \ln(A)$

$\ln\left(\frac{k_1}{k_2} \right) = \frac{E_a}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$

$\text{pH} = \text{pK}_a + \log\left(\frac{[\text{conjugate base}]}{[\text{conjugate acid}]} \right)$

$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{nF} \ln Q$

$E_{\text{cell}}^\circ = \frac{RT}{nF} \ln K^\circ$

$K^\circ = e^{(nF/RT) E_{\text{cell}}^\circ}$

$F = 96485 \text{ J}/\text{V}\cdot\text{mol of electrons}$

$\Delta G^\circ = \Delta H^\circ_{\text{system}} - T\Delta S^\circ_{\text{system}}$

$\Delta G^\circ = -nFE_{\text{cell}}^\circ = -RT \ln K^\circ$

$\Delta G = \Delta G^\circ + RT \ln Q$

$F = 96485 \text{ C}/\text{mol electrons}$

$1A = 1 \text{ C} / \text{sec}$

Quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

1 H 1.0079																	2 He 4.0026				
3 Li 6.941	4 Be 9.0122															5 B 10.811	6 C 12.011	7 N 14.007	8 O 15.999	9 F 18.998	10 Ne 20.180
11 Na 22.990	12 Mg 24.305															13 Al 26.982	14 Si 28.086	15 P 30.974	16 S 32.066	17 Cl 35.453	18 Ar 39.948
19 K 39.098	20 Ca 40.078	21 Sc 44.956	22 Ti 47.88	23 V 50.942	24 Cr 51.996	25 Mn 54.938	26 Fe 55.847	27 Co 58.933	28 Ni 58.69	29 Cu 63.546	30 Zn 65.39	31 Ga 69.723	32 Ge 72.61	33 As 74.922	34 Se 78.96	35 Br 79.904	36 Kr 83.80				
37 Rb 85.468	38 Sr 87.62	39 Y 88.906	40 Zr 91.224	41 Nb 92.906	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.76	52 Te 127.60	53 I 126.90	54 Xe 131.29				
55 Cs 132.91	56 Ba 137.33	57 La 138.91	72 Hf 178.49	73 Ta 180.95	74 W 183.84	75 Re 186.21	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.97	80 Hg 200.59	81 Tl 204.38	82 Pb 207.2	83 Bi 208.98	84 Po (209)	85 At (210)	86 Rn (222)				
87 Fr (223)	88 Ra 226.03	89 Ac 227.03	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs (265)	109 Mt (266)	110 (269)	111 (272)	112 (277)	114		116							

58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.97	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.94	70 Yb 173.04	71 Lu 174.97
90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np 237.05	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (258)	101 Md (258)	102 No (259)	103 Lr (260)

Multiple Choice (3pts each): Circle the letter of the most correct response.

- Rank the 3 states of matter from highest kinetic energy to lowest kinetic energy.
 - Solid, liquid, gas
 - Solid, gas, liquid
 - Gas, solid, liquid
 - Liquid, gas, solid
 - Gas, liquid, solid
- When dissolving a solid in a liquid:
 - Energy is released (exothermic) by breaking solvent-solvent and solute-solute interactions
 - The enthalpy of solution is always positive
 - Formation of solvent-solute interactions is endothermic
 - The boiling point of the solution will be higher than that of the pure solvent
 - The freezing point of the solution will be higher than that of the pure solvent
- Which of the following statements is most correct about colligative properties of an ideal solution?
 - The presence of a solute lowers the boiling point of a solution.
 - The presence of a solute raises the vapor pressure of a solution.
 - Colligative properties depend upon the number of solute particles, not on the identity of the solute particles.
 - The presence of a solute raises the freezing point of a solution.
 - These statements are all correct.
- The volume of a gas:
 - Decreases as the temperature increases
 - Remains constant as the amount of gas is increased
 - Is always a constant
 - Decreases as the pressure increases
 - Increases as the kinetic energy increases
- Under which of the following conditions is a gas *least* “ideal”?
 - Room temperature, 25°C
 - High temperature, high pressure
 - Low temperature, high pressure
 - Low temperature, low pressure
 - High temperature, low pressure
- Which of the following is *not* a correct gas law relationship?
 - $PV = nRT$
 - $n_1T_1 = n_2T_2$
 - $V_1n_1 = V_2n_2$
 - $P_1V_1 = P_2V_2$
 - $P_1 / T_1 = P_2 / T_2$
- Carbon tetrabromide (CBr₄) has a higher boiling point than carbon tetrafluoride (CF₄) because:
 - The bonds in CF₄ are polar but the bonds in CBr₄ are not
 - CBr₄ has a higher molecular weight than CF₄
 - CF₄ is a polar molecule but CBr₄ is not
 - CF₄ has stronger intermolecular forces than CBr₄
 - CF₄ is a gas at room temperature

8. What is the volume of 2.834mols of ideal gas at 11.14°C and 4.64atm pressure? (8pts)

Answer 8:

9. You have a 82.91L sample of gas at 21.61°C. What is the volume of this gas if the temperature is increased to 62.93°C? (8pts)

Answer 9:

10. You have prepared a solution by dissolving 15.443g of magnesium bromide in enough water to make 250.0mL of solution. What is the *molarity* of this solution? (8pts)

Answer 10:

11. You have prepared a solution by dissolving 17.228g of ammonium chlorate in 100.0g of water. What is the *molality* of this solution? (8pts)

Answer 11:

12. You have prepared a solution by diluting 25.00mL of a 1.885M aqueous solution of iron(II) sulfate to a total volume of 125.0mL. What is the *molarity of iron(II)* of this solution? (8pts)

Answer 12:

13. What is the boiling point of a solution made by dissolving 13.861g of nickel(II) nitrate in 150.0g of water? (12pts)

Answer 13:


14. Each of the following solids is dissolved in separate beakers containing 500.0mL of water. Rank the solutions from lowest freezing point to highest freezing and explain your answer. (12pts)
- 0.5mols sodium phosphate
 - 0.5mols calcium nitrate
 - 0.6mols magnesium chloride
 - 0.6mols ammonium carbonate

15. A laboratory technician prepares a solution by weighing out 39.225g of potassium bromide and dissolving it in enough water to make 150.00mL of solution. The technician labels the solution “1.3M KBr(aq)”. Why is this not correct? What should the laboratory technician do to correct the error? Calculate a correct concentration for this solution. (8pts)

16. How much energy is required to heat 500.0g of water from -37.25°C to 21.39°C ? $\{C_s(\text{ice}) = 2.09 \text{ J/g}\cdot\text{K}; C_s(\text{water}) = 4.184 \text{ J/g}\cdot\text{K}; C_s(\text{steam}) = 2.01 \text{ J/g}\cdot\text{K}; \Delta H_{\text{fusion}}(\text{water}) = 6.02 \text{ kJ/mol}; \Delta H_{\text{vaporization}}(\text{water}) = 40.7 \text{ kJ/mol}\}$ (15pts)

Answer 16:

17. Some salts have enough covalent bond character that they do not completely dissociate when dissolved in water. You have performed an experiment in which you have made a solution by dissolving 29.531g of tin(IV) bromide in 250.00mL of water. The observed freezing point of this solution is -1.50°C . Which of the following equations is most consistent with your observed freezing point? Explain your answer with explicit calculations. (15pts)



18. You have reacted 8.042g of aluminum metal with 150.0mL of 4.623M HCl(aq) to produce hydrogen gas and aluminum chloride. How many liters of hydrogen gas can be produced at 41.61°C and 1.137atm? (15pts)

Answer 18:

19. A newly discovered protein has been isolated from seeds of a tropical plant and needs to be characterized. A total of 0.137g of this protein was dissolved in enough water to produce 2.00mL of solution. At 31.68°C the osmotic pressure produced by the solution was 0.134atm. What is the molar mass of the protein? (15pts)

Answer 19: