

Chemistry 210

Exam 4

Be sure to put your name on each page. This page can be removed from your exam so that you will have a Periodic Table handy throughout the exam, it does not need to be turned in. Show all your work for problems which require any sort of calculation, no credit will be given for answers without work shown. If you have shown a significant amount of work or multiple drawings for a problem, draw a box around what you consider your final answer.

Avogadro's Number = 6.022×10^{23} units/mol

$32.00^\circ\text{F} = 0.000^\circ\text{C} = 273.15\text{K}$

Density of Water = $1.000 \frac{\text{g}}{\text{mL}}$

$R = 0.08206 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}} = 8.314 \frac{\text{J}}{\text{mol}\cdot\text{K}}$

$1\text{atm} = 760\text{torr} = 760\text{mmHg} = 101.325\text{kPa}$

$PV = nRT$

$\Delta T_{\text{fp/bp}} = k_{\text{fp/bp}} \cdot m \cdot i$

For water: $k_{\text{fp}} = -1.86^\circ\text{C}/m$
 $k_{\text{bp}} = 0.512^\circ\text{C}/m$

$P_1 = X_1 P_1^\circ$

$\Pi = MRTi$

$C_1 V_1 = C_2 V_2$

$E = C_s \cdot g \cdot \Delta T$

Quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Integrated Rate Laws:

0th $[A]_t = -kt + [A]_0$

1st $\ln[A]_t = -kt + \ln[A]_0$

2nd $1/[A]_t = kt + 1/[A]_0$

$k = Ae^{-E_a/RT}$

$$\ln(k) = \left(\frac{-E_a}{R} \right) \left(\frac{1}{T} \right) + \ln(A)$$

$$\ln\left(\frac{k_1}{k_2} \right) = \frac{E_a}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

$$\text{pH} = \text{pK}_a + \log\left(\frac{[\text{conjugate base}]}{[\text{conjugate acid}]} \right)$$

$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{nF} \ln Q$

$E_{\text{cell}}^\circ = \frac{RT}{nF} \ln K^\circ$

$K^\circ = e^{(nF/RT) E_{\text{cell}}^\circ}$

$F = 96485 \frac{\text{J}}{\text{V}\cdot\text{mol of electrons}}$

$\Delta G^\circ = \Delta H^\circ_{\text{system}} - T\Delta S^\circ_{\text{system}}$

$\Delta G^\circ = -nFE_{\text{cell}}^\circ = -RT \ln K^\circ$

$\Delta G = \Delta G^\circ + RT \ln Q$

$F = 96485 \frac{\text{C}}{\text{mol electrons}}$

$1A = 1C / \text{sec}$


1 H 1.0079																	2 He 4.0026				
3 Li 6.941	4 Be 9.0122															5 B 10.811	6 C 12.011	7 N 14.007	8 O 15.999	9 F 18.998	10 Ne 20.180
11 Na 22.990	12 Mg 24.305															13 Al 26.982	14 Si 28.086	15 P 30.974	16 S 32.066	17 Cl 35.453	18 Ar 39.948
19 K 39.098	20 Ca 40.078	21 Sc 44.956	22 Ti 47.88	23 V 50.942	24 Cr 51.996	25 Mn 54.938	26 Fe 55.847	27 Co 58.933	28 Ni 58.69	29 Cu 63.546	30 Zn 65.39	31 Ga 69.723	32 Ge 72.61	33 As 74.922	34 Se 78.96	35 Br 79.904	36 Kr 83.80				
37 Rb 85.468	38 Sr 87.62	39 Y 88.906	40 Zr 91.224	41 Nb 92.906	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.76	52 Te 127.60	53 I 126.90	54 Xe 131.29				
55 Cs 132.91	56 Ba 137.33	57 La 138.91	72 Hf 178.49	73 Ta 180.95	74 W 183.84	75 Re 186.21	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.97	80 Hg 200.59	81 Tl 204.38	82 Pb 207.2	83 Bi 208.98	84 Po (209)	85 At (210)	86 Rn (222)				
87 Fr (223)	88 Ra 226.03	89 Ac 227.03	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs (265)	109 Mt (266)	110 (269)	111 (272)	112 (277)	114		116							

58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm (145)	62 Sm 150.36	63 Eu 151.97	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.94	70 Yb 173.04	71 Lu 174.97
90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np 237.05	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (258)	101 Md (258)	102 No (259)	103 Lr (260)



Multiple Choice (4pts each): Circle the letter of the most correct response.

- A large negative change in free energy means:
 - The reaction is very slow
 - The reaction is exothermic
 - The reaction is not spontaneous
 - The system is becoming more disordered
 - The reaction is spontaneous
- A reaction will be spontaneous at relatively high temperature and non-spontaneous at relatively low temperature if:
 - $\Delta H^\circ_{\text{system}} > 0$ and $\Delta S^\circ_{\text{system}} > 0$
 - $\Delta H^\circ_{\text{system}} < 0$ and $\Delta S^\circ_{\text{system}} > 0$
 - $\Delta H^\circ_{\text{system}} > 0$ and $\Delta S^\circ_{\text{system}} = 0$
 - $\Delta H^\circ_{\text{system}} > 0$ and $\Delta S^\circ_{\text{system}} < 0$
 - $\Delta H^\circ_{\text{system}} < 0$ and $\Delta S^\circ_{\text{system}} < 0$
- For a reaction with a small negative ΔS :
 - Heat is liberated by the reaction
 - The system is becoming more ordered
 - The reaction is not spontaneous
 - The disorder of the system is increasing
 - The reaction proceeds very quickly
- If the change in enthalpy for a reaction is positive and the change in entropy is negative:
 - The system is becoming more disordered
 - The reaction releases heat
 - The reaction will be spontaneous at all temperatures
 - The reaction will be non-spontaneous at all temperatures
 - The reaction will be spontaneous only at low temperatures
- A reaction will be product-favored/spontaneous if:
 - $\Delta G^\circ < 0$
 - $K_{\text{eq}} < 1$
 - $\Delta H > 0$
 - $\Delta S^\circ < 0$
 - $E^\circ_{\text{cell}} < 0$
- How are the change in Gibbs' Free Energy and the equilibrium constant for a reaction related?
 - As K approaches zero, ΔG approaches zero
 - The value of ΔG is equal to $(-\log K)$
 - As ΔG gets more positive, K approaches 1
 - They're not.
 - As ΔG gets more negative, K gets very large
- In a spontaneous electrochemical voltaic cell, which of the following is *true*?
 - The cell potential is zero
 - Oxidation occurs at the cathode
 - Electrons flow from the cathode to the anode
 - Cations flow through the salt bridge from the cathode to the anode
 - The metal cathode gains mass as the cell reaction proceeds

8. For a spontaneous redox reaction, which of the following is *false*?
- Oxidation is the process of losing electrons
 - Gaining electrons is reduction
 - Electrons appear on the left side of the oxidation half reaction
 - Water molecules are added to balance any extra oxygen atoms
 - ΔG is negative.
9. Give the oxidation number for each atom in the following formulas. (15pts)




10. For each of the following reactions, predict whether the sign of ΔS° will be positive or negative and explain your answer. (15pts)



11. For liquid water $S^\circ = +69.91 \text{ J/mol}\cdot\text{K}$ and for solid water $S^\circ = +41.3 \text{ J/mol}\cdot\text{K}$. Explain this difference. (10pts)

12. You are studying a process for which $\Delta H^\circ = -28.38 \text{ kJ/mol}$ and $\Delta S^\circ = -137.62 \text{ J/mol}\cdot\text{K}$. What is ΔG° for this process at 25.00°C ? Will the reaction be more or less spontaneous at 20.00°C ? (10pts)
13. Ammonia $\{\text{NH}_3(\text{g})\}$ can burn in oxygen to form nitrogen dioxide and water. How much {Gibb's Free} energy can be liberated by burning 23.948g of ammonia in an unlimited supply of oxygen? (10pts)
14. You are studying the reaction of propanoic acid $\{\text{C}_2\text{H}_5\text{CO}_2\text{H}(\text{l})\}$ with ethyl amine $\{\text{C}_2\text{H}_5\text{NH}_2(\text{l})\}$ to produce N-ethylpropanamide $\{\text{C}_5\text{H}_{11}\text{NO}(\text{l})\}$ and water. When you run the reaction at 23.61°C , you find that ΔG for this reaction is -18.62 kJ/mol and $\Delta S = +61.37 \text{ J/mol}\cdot\text{K}$ (12pts)
- Is the reaction endothermic or exothermic? (*Explain your answer with explicit calculations.*)
 - Over what temperature range is this reaction spontaneous?
15. You have burned 72.612g of ethane $\{\text{C}_2\text{H}_6(\text{g})\}$ in oxygen to form carbon dioxide and water. If all of the Gibb's Free Energy liberated by this reaction is used to decompose iron(III) chloride to iron metal and chlorine gas, how many grams of iron metal will be formed? (16pts)

16. For each of the following pairs of half-reactions/half-cells, determine the voltage of the spontaneous reaction/cell and write a balanced equation for the spontaneous reaction, identifying the oxidation and reduction half-reactions. (10pts each)



Thermodynamic Values at 25°C:

Substance	ΔH_f° (kJ/mol)	S° (J/mol·K)	ΔG_f° (kJ/mol)
$\text{NH}_3(\text{g})$	-46.11	+192.45	-16.45
$\text{O}_2(\text{g})$	0	+205.138	0
$\text{NO}_2(\text{g})$	+33.18	+240.06	+51.31
$\text{H}_2\text{O}(\text{g})$	-241.8	+188.8	-228.6
$\text{H}_2\text{O}(\text{l})$	-285.8	+69.91	-237.2
$\text{C}_2\text{H}_6(\text{g})$	-84.68	+229.2	-32.0
$\text{CO}_2(\text{g})$	-393.5	+213.6	-394.4
$\text{FeCl}_3(\text{s})$	-400.39	+142.3	-334.18
$\text{Fe}(\text{s})$	0	+27.15	0
$\text{Cl}_2(\text{g})$	0	+222.96	0

Standard Reduction Potentials at 25°C:

Half cell	E°_{red} (volts)	Half cell	E°_{red} (volts)
$\text{Al}^{3+}(\text{aq}) \text{Al}(\text{s})$	-1.676	$\text{Br}_2(\text{g}) \text{Br}^{-1}(\text{aq})$	+1.06
$\text{Pb}^{4+}(\text{aq}) \text{Pb}^{2+}(\text{aq})$	+1.69	$\text{NO}_3^{-1}(\text{aq}) \text{NO}(\text{g})$	+0.96
$\text{GeO}_2(\text{s}) \text{GeO}(\text{s})$	-0.370	$\text{Cr}_2\text{O}_7^{-2}(\text{aq}) \text{Cr}^{3+}(\text{aq})$	+1.33